



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Book

Search PubMed  for  Go Clear  
Limits Preview/Index History Clipboard Details

Display Abstract  Show: 20  Sort  Send to Text

1: Gene 1999 May 17;232(1):43-51

[Related Articles, Links](#)

[LIFE SCIENCE](#)  
[FULL-TEXT ARTICLE](#)

### Characterization of the biotin biosynthesis pathway in *Saccharomyces cerevisiae* and evidence for a cluster containing BIO5, a novel gene involved in vitamers uptake.

Phalip V, Kuhn I, Lemoine Y, Jeltsch JM.

Laboratoire de Biotechnologie, Ecole Supérieure de Biotechnologie de  
Strasbourg, Boulevard Sebastien Brandt, 67400, Illkirch-Graffenstaden, France.  
vincent.phalip@esbs.u-strasbg.fr

An engineered mutant of *Saccharomyces cerevisiae* affected in biotin biosynthesis has been isolated. This mutant allowed the characterization of a bi cluster (BIO3-4-5). We demonstrate that BIO3 (YNR058w) and BIO4 (YNR057c) encode, respectively, a 7, 8-diaminopelargonic acid aminotransferase and a dethiobiotin synthase, involved in the biotin biosynthesis pathway. A novel gene, BIO5 (YNR056c), is present immediately downstream from BIO4. This gene encodes Bio5p, a protein with 11 putative transmembrane regions. Uptake experiments performed with labeled 7-keto 8-aminopelargonic acid indicate that Bio5p is responsible for transport into the cell of 7-keto 8-aminopelargonic acid.

PMID: 10333520 [PubMed - indexed for MEDLINE]

Display Abstract  Show: 20  Sort  Send to Text

Write to the Help Desk  
NCBI | NLM | NIH  
Department of Health & Human Services  
[Freedom of Information Act](#) | [Disclaimer](#)